

Original Effective Date: 02/28/2008 Current Effective Date: 01/07/2023 Last P&T Approval/Version: 10/26/2022

Next Review Due By: 10/2023 Policy Number: C14676-A

## Hyaluronic Acid Injections\_Viscosupplementation for Knee OA

## **PRODUCTS AFFECTED**

Durolane (sodium hyaluronate injection), Euflexxa (sodium hyaluronate injection), Gel-One (sodium hyaluronate injection), Gelsyn-3 (sodium hyaluronate injection), GenVisc 850 (sodium hyaluronate injection), Hyalgan (sodium hyaluronate injection), Hymovis (high molecular weight viscoelastic hyaluronan injection), Monovisc (high molecular weight hyaluronan injection), Orthovisc (high molecular weight hyaluronan injection), Supartz (sodium hyaluronate), Synvisc (hylan G-F 20 sodium hyaluronate injection), Synvisc-One (hylan G-F 20 sodium hyaluronate injection), TriVisc (sodium hyaluronate injection), Visco-3 (sodium hyaluronate injection), Synojoynt (sodium hyaluronate injection), Triluron (sodium hyaluronate injection), 1% Sodium Hyaluronate

## **COVERAGE POLICY**

Coverage for services, procedures, medical devices, and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines.

### **Documentation Requirements:**

Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive.

#### **DIAGNOSIS:**

Osteoarthritis (OA) of the Knee

## **REQUIRED MEDICAL INFORMATION:**

This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. If a drug within this policy receives an updated FDA label within the last 180 days, medical necessity for the member will be reviewed using the updated FDA label information along with state and federal requirements, benefit being administered and formulary preferencing. Coverage will be determined on a case-by case basis until the criteria can be updated through Molina Healthcare, Inc. clinical governance. Additional information may be required on a case-by-case basis to allow for

adequate review, Molina Healthcare, Inc. confidential and proprietary © 2022

This document contains confidential and proprietary information of Molina Healthcare and cannot be reproduced, distributed, or printed without written permission from Molina Healthcare. This page contains prescription brand name drugs that are trademarks or registered trademarks of pharmaceutical manufacturers that are not affiliated with Molina Healthcare.

Page 1 of 9

## A. OSTEOARTHRITIS (OA) OF THE KNEE

- Diagnosis of osteoarthritis of the knee AND
- Documentation of member's affected knee(s): Left, right or both knees to be treated.
   NOTE: Bilateral injections may be allowed only if both knees meet criteria.
   AND
- 3. Prescriber attests that member has no evidence of inflammatory arthritis (e.g., rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis, juvenile idiopathic arthritis, and systemic lupus erythematosus) and other causes of musculoskeletal pain, including referred pain, bursitis, and inflammatory rheumatic diseases have been ruled out.

  Informational Note: Safety and efficacy in joints with severe inflammation have not been established. There are no studies that have evaluated the efficacy of hyaluronate derivatives in patients with OA and coexistent other inflammatory conditions such as rheumatoid arthritis. AND
- 4. Prescriber attests that member has NOT had a history of failure on hyaluronic intraarticular injection (applicable to member's medical history prior to, and as a Molina member). Requests will not be authorized for members who have failed to respond to ANY previous viscosupplementation therapy. Informational Note: There is a lack of reliable evidence that any one brand of viscosupplement is more effective to other brands for medically necessary indications. There are also a lack of studies demonstrating that individuals who fail to respond to one brand of viscosupplement will respond to other brands of viscosupplements. AND
- 5. Prescriber attests that surgical knee replacement is not a planned treatment option within 6 months of intra-articular viscosupplementation administration

  Informational Note: There is no data to suggest efficacy of hyaluronate derivatives in patients who have had total knee arthroplasty in the targeted knee.

  AND
- 6. Documentation that the member has tried and has experienced Ineffectiveness/failure (defined as symptoms inadequately controlled after an adherent 3-month trial unless specified), clinical intolerance (defined as an intolerance to the drug or its excipients); or FDA-labeled contraindication(s)\* in the affected joint to TWO of the three following modalities of therapy [DOCUMENTATION REQUIRED]
  - At least one course of physical therapy (PT) for knee osteoarthritis
     OR
  - b. At least TWO of the following pharmacologic therapies [verification of therapies required]:
    - Non-steroidal anti-inflammatory drugs [Contraindications may include:
      - 1) Compromised GI function or at risk of GI bleeding due to the adverse events of NSAIDs, 2) Concomitant anticoagulant therapy for any condition,
      - 3) Cardiovascular or renal risk factors precluding use of COX-2 inhibitors]
    - ii. Acetaminophen (up to 1 g 4 times/day)
    - iii. Tramadol (Ultram/XR, generics)
    - iv. Duloxetine (Cymbalta, generics)

OR

 At least TWO injections of IA corticosteroids to the affected knee [Contraindications may include: increase in risk of local or systemic bacterial infection]

Molina staff: Verify pharmacy claims data for above medications and compliance. For new members to Molina Healthcare, confirm medications use in medical or chart notes. Non- compliance or non-adherence does <u>not</u> constitute therapeutic failure.

AND

7. Documentation of prescriber baseline pain and functional capacity assessment by any type of

objective/quantification method, such as Visual Analog Scale for pain, joint mobility, effusion, and/or patient-response-based questionnaires [DOCUMENTATION REQUIRED]

AND

 IF THIS IS A PHARMACY BENEFIT REQUEST FOR A NON-FORMULARY/NON-PREFERRED PRODUCT: Documentation of CLINICAL MEDICAL rationale for the inability to utilize the preferred formulary hyaluronate products. OR

FOR MEDICAL BENEFIT CLAIMS (i.e., HCPC billing by a prescriber) REQUESTS: Documentation of CLINICAL MEDICAL rationale for the inability to utilize the preferred hyaluronate product.

#### CONTINUATION OF THERAPY:

## A. OSTEOARTHRITIS (OA) OF THE KNEE

- 1. Documentation of positive response to therapy documented by ALL of the following [DOCUMENTATION REQUIRED:
  - a. Significant improvement in pain and functional capacity as the result of the previous series of injections as indicated by any type of objective/quantification method, such as Visual Analog Scale for pain, joint mobility, reduction in effusion, and/or patient-response-based questionnaires

NOTE: If the initial or prior series of injections is not proven, or documented as beneficial to the member, it is not considered medically necessary to repeat the therapy and a repeat series of injections will not be authorized.

AND

 A significant reduction in the dose/utilization of pharmacologic therapy (NSAIDs or other analgesics or anti- inflammatory medication)
 OR

A reduction in the number of accompanying intra-articular corticosteroid injections during the six month period following the previous series of injections

**AND** 

- At least six months have elapsed since the initial or prior treatment cycle AND
- 3. Prescriber attests member has not had total or partial joint replacement surgery Informational Note: There are no clinical trials evaluating the use of sodium hyaluronate in persons following total or partial joint replacement surgery.

## **DURATION OF APPROVAL:**

Initial authorization: A single course per joint every 6 months

Continuation of treatment: At least **six (6) months** have elapsed since the initial or prior treatment cycle for the respective joint. Re-authorization is required every **6 months** to determine effectiveness of therapy and continued need based on documented positive clinical response. *Refer to 'Continuation of Therapy' section.* 

#### PRESCRIBER REQUIREMENTS:

Prescribed by, or in consultation with, a board-certified orthopedic surgeon, pain management specialist, rheumatologist, physical medicine and rehabilitation specialist, or a sports medicine specialist. [If prescribed in consultation, consultation notes must be submitted with initial request and reauthorization

## **AGE RESTRICTIONS:**

18 years of age and older

## **QUANTITY:**

| Table 1: FDA Labeled Dosage per Treatment Course per<br>Joint |  |                   |  |
|---|--|-------------------|--|
| Drug  | Dose   | Total Injections  |  |
| Durolane (hyaluronic acid)                                    | 60 mg (3 ml) intra-articularly <b>once</b>   | 1 injection       |  |
| Euflexxa (1% sodium hyaluronate)                              | 20 mg (2 mL) intra-articularly once weekly for 3 weeks 3 injection   |                   |  |
| Gel-One (Cross-linked<br>Hyaluronate)                         | 30 mg (3 mL) intra-articularly <b>once</b>   | 1 injection       |  |
| Gelsyn-3 (0.84%<br>sodium hyaluronate)                        | 16.8 mg (2 mL) intra-articularly once weekly for 3 weeks 3 inje  |                   |  |
| GenVisc 850<br>(sodium hyaluronate)                           | 5 mg (2.5 mL) intra-articularly once weekly for 5 veeks. Some patients may benefit from 3 injections 3 to 5 injection iven at weekly intervals |                   |  |
| Hyalgan (sodium<br>hyaluronate)                               | 20 mg (2 mL) intra-articularly once weekly for 5 weeks.<br>Some patients may benefit with 3 injections given at<br>weekly intervals            | 3 to 5 injections |  |
| Hymovis (high molecular weight hyaluronan)                    | 24 mg (3 mL) intra-articularly once weekly for 2 weeks 2 injection   |                   |  |
| Monovisc (high molecular weight hyaluronan)                   | 88 mg (4 mL) intra-articularly once  | 1 injection       |  |
| Orthovisc (high molecular weight hyaluronan)                  | 30 mg (2 mL) intra-articularly once weekly for 3 or 4 weeks  | 3 to 4 injections |  |
| 1% Sodium Hyaluronate   | 20 mg (2 mL) intra-articularly once weekly for 3 weeks   | 3 injections      |  |
| Supartz FX  | 25 mg (2.5 mL) intra-articularly once weekly for 5 weeks. Supartz FX Some patients may benefit with 3 injections given at weekly intervals     |                   |  |
| SynoJoynt (sodium<br>hyaluronate                              | 20 mg (2 mL) intra-articularly once weekly for 3 weeks 3 injections  |                   |  |
| Synvisc One (Hylan G-F<br>20)                                 | 48 mg (6 mL) intra-articularly once 1 injection  |                   |  |
| Synvisc (Hylan G-F 20)  | 16 mg (2 mL) intra-articularly once weekly for 3 weeks   | 3 injections      |  |
| Triluron (sodium<br>hyaluronate)                              | 20 mg (2 mL) once weekly for 3 weeks   | 3 injections      |  |
| TriVisc (sodium hyaluronate)                                  | 25 mg (2.5 mL) once weekly for 3 weeks   | 3 injections      |  |
| Visco-3   | 25 mg (2.5 mL) intra-articularly once weekly for 3 weeks   | 3 injections      |  |

## Maximum Quantity Limits - Based on FDA label

#### PLACE OF ADMINISTRATION:

The recommendation is that intra-articular injection medications in this policy will be for pharmacy or medical benefit coverage and be administered in a place of service that is a non-hospital facility-based location.

## **DRUG INFORMATION**

## **ROUTE OF ADMINISTRATION:**

Intra-articular injection directly into the knee joint

#### DRUG CLASS:

Antirheumatic Miscellaneous

#### FDA-APPROVED USES:

Osteoarthritis of the Knee

Treatment of pain in osteoarthritis of the knee in patients who have failed nonpharmacologic treatment and simple analgesics

(*Durolane*, *Euflexxa*, *Gel-One*, *Gelsyn-3*, *GenVisc* 850, *Hyalgan*, *Hymovis*, *Monovisc*, *Orthovisc*, sodium hyaluronate [Teva], *Supartz FX*, *Synvisc-One*, *Triluron*, *TriVisc*, *Visco-3*) or nonsteroidal anti-inflammatory drugs (*Gel-One*)

## **COMPENDIAL APPROVED OFF-LABELED USES:**

None

## **APPENDIX**

#### **APPENDIX:**

Comparison of OA Management Guidelines

- Comparison of 2 guidelines (AAOS 2013, VA/DoD 2014) on nonsurgical management of osteoarthritis of the knee Reference: <u>National Guideline Clearinghouse 2016 Jun 13:50210</u>
- Comparison of 16 guidelines on the management of osteoarthritis Reference:
   Semin Arthritis Rheum 2014 Jun;43(6):701

## **BACKGROUND AND OTHER CONSIDERATIONS**

### BACKGROUND:

Clinical studies of sodium hyaluronate and hylan G-F-20 have demonstrated that injection of these agents into the joint space of osteoarthritic knees is sometimes marginally more effective than placebo procedures in reduction of pain and improvement in functional capacity in some patients. These marginal beneficial results are more pronounced with the larger molecular weight compound hylan G-F20.

There is no data indicating that these agents reverse or delay the osteoarthritic process in the injected joints. The long-term effects of repeated injections are unknown.

## **Clinical Practice Guidelines**

• The majority of guidelines did not find sufficient evidence to make a recommendation for or against the use of HA for knee OA. Refer to 'Supplemental Information' section for additional

references and links for comparisons of guidelines.

- There is inconsistent evidence and limited effectiveness data that viscosupplementation, or HA
  products, produces clinically relevant improvements in pain and functioning for OA of the knee
  and no evidence to suggest it delays the progression of OA nor the progression to knee
  replacement.
- Several major practice guidelines have been unable to recommend intraarticular HA, with others recommending against its use with several other major organizations.

## **National and Specialty Organizations**

American Academy of Orthopedic Surgeons (AAOS) In the second edition of the evidence-based guidelines on treatment of OA of the knee, the AAOS issued a "Strong" recommendation against the use of HA for knee OA due to lack of efficacy (AAOS, 2013).

National Institute for Health and Care Excellence (NICE 2014) recommended against HA for knee OA

American College of Rheumatology (ACR 2012) clinical practice guidelines on osteoarthritis indicate they have no recommendations regarding the use of IAHA in the knee. Recommendations for the use of pharmacologic therapies in knee OA include acetaminophen, oral and topical NSAIDs, tramadol and intra- articular corticosteroid injections. The ACR conditionally recommends IA-HA for patients who had inadequate relief with initial treatment for knee OA in evidence-based guidelines (Hochberg et al., 2012). An update of this guideline is anticipated in 2018 (Osteoarthritis: Call for Public Comment on Project Plan).

American Medical Society for Sports Medicine (AMSSM) Based on findings from their systematic review with network meta-analysis, the AMSSM recommended IA-HA for appropriate patients with knee OA. Criteriafor appropriate patients were not reported. (Trojian et al., 2016)

Osteoarthritis Research Society International (OARSI) 2014 guideline update provided an "uncertain" ommendation for IAHA, indicating an overall small effect size on pain, inconsistent results among the available meta-analyses, and one meta-analysis signaling potential for serious safety concerns, influenced their recommendation.

**ECRI Institute. Viscosupplementation for Treating Osteoarthritic Knee Pain** A 2019 ECRI report on viscosupplementation summarized evidence from 8 systematic reviews and 6 RCTs (total patients = 12,775) to be inconclusive for treating knee pain due to OA. While IA HA injections may provide relief in some patients, uncertainty remains about the most effective formulations, which populations benefit most, and whether HA should be combined with other agents to increase efficacy. (ECRI; 2019)

## CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:

The following are considered contraindications/exclusions based on insufficient evidence:-

- 1. Hypersensitivity to hyaluronan preparations, or allergies to avian or avian-derived products (including eggs, feathers, or poultry) NOTE: Not applicable to Orthovisc Informational Note: HA is derived from chicken combs (Gel-One, Hyalgan, Supartz FX, Synvisc, Synvisc-One, Visco-3) or bacterial cells (Euflexxa, Gelsyn-3, Hymovis, Monovisc, Orthovisc)
- 2. Hymovis, Monovisc and Orthovisc only: known hypersensitivity to gram positive bacterial proteins
- 3. Monovisc and Durolane only: known systemic bleeding disorders
- 4. Any of the following conditions:
  - a. Active inflammatory joint disease or synovitis affecting the knee, such as crystal induced synovitis, rheumatoid arthritis
  - b. Knee joint infections (septic arthritis) or a local skin disease or infection in the area of the injection site

# Drug and Biologic Coverage Criteria c. Pregnant or nursing

The following are considered conditions for **discontinuation of treatment** and re-treatment may not be authorized:-

- 1. Intolerable adverse effects or absence of unacceptable toxicity from the drug
- 2. Persistent and uncorrectable problems with adherence to treatment
- 3. Poor response to treatment as evidenced by physical findings and/or clinical symptoms

The following are considered **experimental**, **investigational and unproven** based on insufficient evidence:

1. Any indications other than those listed above (i.e., Injection of these products for indications other than the diagnosis OA, or use of the requested products for injection into any joint other than the knee)

#### COSMETIC USE IS NOT A COVERED BENEFIT

The FDA has approved several products containing a transparent HA gel to improve the contours of the skin. These products are used to treat acne, scars and wrinkles on the skin by temporarily adding volume to facial tissue and restoring a smoother appearance to the face (may not be an all-inclusive list):

- Restylane injectable gel received Premarket approval (PMA) approval March 25, 2005
- Perlane injectable gel received PMA approval May 2, 2007
- Hylaform received PMA approval April 22, 2004
- Juvéderm 24HV, Juvéderm 30 & Juvéderm 30HV Gel Implants received PMA approval June 2, 2006

#### **OTHER SPECIAL CONSIDERATIONS:**

None

#### **CODING/BILLING INFORMATION**

Note: 1) This list of codes may not be all-inclusive. 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement

| HCPCS | Description  |
|-------|--|
| J7318 | Hyaluronan or derivative, durolane, for intra-articular injection, 1 mg  |
| J7320 | Hyaluronan or derivative, GenVisc 850, for intra-articular injection, 1 mg   |
| J7321 | Hyaluronan or derivative, hyalgan, supartz or visco-3, for intra-articular injection, per dose Effective Date: 04/01/2021  |
| J7322 | Hyaluronan or derivative, Hymovis, for intra-articular injection, 1 mg   |
| J7323 | Hyaluronan or derivative, Euflexxa, for intra-articular injection, per dose  |
| J7324 | Hyaluronan or derivative, Orthovisc, for intra-articular injection, per dose   |
| J7325 | Hyaluronan or derivative, Synvisc or Synvisc-One, for intra- articular injection, 1 mg   |
| J7326 | Hyaluronan or derivative, Gel-One, for intra-articular injection, per dose   |
| J7327 | Hyaluronan or derivative, Monovisc, for intra-articular injection, per dose  |
| J7328 | Hyaluronan or derivative, GelSyn-3, for intra-articular injection, 0.1 mg  |
| J7329 | Hyaluronan or derivative, trivisc, for intra-articular injection, 1 mg   |
| J7331 | Hyaluronan or derivative, synojoynt, for intra-articular injection, 1 mg   |
| J7332 | Hyaluronan or derivative, triluron, for intra-articular injection, 1 mg  |
| J7333 | Hyaluronan or derivative, VISCO-3, for intra-articular injection, per doseEffective July 1, 2020 Discontinue existing HCPCS code J7333 "Hyaluronan or derivative, visco-3, for intra-articular injection, per dose."Effective Date: 03/31/2021 |

## **AVAILABLE DOSAGE FORMS:**

Several HA agents are available, with varying molecular weights and injections per course of treatment (single injection HAs and those requiring 3 to 5 injections per course of treatment). Refer to Table 1 in 'Coverage Policy' section above

## **REFERENCES**

- 1. Centers for Medicare & Medicaid Services (CMS). Medicare Coverage Database. National Coverage Determination (NCD) Search. Accessed at: CMS; CMS NCD
- 2. Centers for Disease Control and Prevention. Osteoarthritis Fact Sheet. Available at https://www.cdc.gov/arthritis/basics/osteoarthritis.htm. February 7, 2018; Accessed February 2021.
- Durolane (sodium hyaluronate) [prescribing information]. Durham, NC: Bioventus LLC; October 2017
- 4. Euflexxa (sodium hyaluronate) [prescribing information]. Parsippany, NJ: Ferring Pharmaceuticals Inc; July 2016.
- 5. Hyalgan (hyaluronic acid derivative) [prescribing information]. Parsippany, NJ: Fidia Pharma; May 2014.
- Hymovis (hyaluronic acid derivative) [prescribing information]. Parsippany, NJ: Fidia Pharma: October 2015.
- 7. Gel-One [prescribing information]. Warsaw, IN: Zimmer; May 2011.
- 8. Gelsyn-3 (sodium hyaluronate) [prescribing information]. Durham, NC: Bioventus; 2016.
- 9. GenVisc 850 (sodium hyaluronate) [prescribing information]. Doylestown, PA: OrthogenRx Inc; no date.
- 10. Monovisc [prescribing information]. Bedford, MA: Anika Therapeutics; December 2013.
- 11. Orthovisc [prescribing information]. Raynham, MA: Anika Therapeutics; June 2005.
- 12. Synvisc [package insert]. Ridgefield, NJ: Genzyme Biosurgery; September 2014.
- 13. Synvisc One [package insert]. Ridgefield, NJ: Genzyme Biosurgery; September 2014.
- 14. Supartz FX (sodium hyaluronate) [prescribing information]. Durham, NC: Bioventus; April 2015.
- 15. Triluron (sodium hyaluronate) [prescribing information]. Florham Park, NJ: Fidia Pharma USA Inc.; July 2019.
- 16. TriVisc (sodium hyaluronate) [prescribing information]. Doylestown, Pennsylvania: OrthogenRx Inc; September 2018.
- 17. Visco-3 (sodium hyaluronate) [prescribing information]. Warsaw, IN: Zimmer; received April 2017.
- 18. Bannuru RR, Schmid CH, Kent DM, et al. Comparative Effectiveness of Pharmacologic Interventions for Knee Osteoarthritis: a systematic review and network meta-analysis. Ann Intern Med. 2015 Jan 6;162(1):46-54. doi: 10.7326/M14-1231. PMID: 25560713. Accessed Mar 2021
- 19. Jüni P, Hari R, Rutjes AW, et al. Intra-articular corticosteroid for knee osteoarthritis. Cochrane Database Syst Rev 2015: CD005328. https://www.cochranelibrary.com/cdsr/doi/10.1002/14651858.CD005328.pub3/epdf/full Accessed Mar 2021
- 20. Agency for Healthcare Research and Quality. Treatment of Osteoarthritis of the Knee: An Update Review. Comparative Effectiveness Review No. 190. (Prepared by the RAND Southern California Evidence-based Practice Center under Contract No. 290-2015-00010-I.) AHRQ Publication No.17-EHC011-EF. May 2017. Available at: Link Accessed Feb 2021
- 21. American Academy of Orthopaedic Surgeons (AAOS). Treatment of Osteoarthritis of the Knee. 2nd Edition. Evidence-Based Clinical Practice Guideline. Available at: AAOS Accessed Feb 2021
- 22. American Medical Society for Sports Medicine (AMSSM). AMSSM scientific statement concerning viscosupplementation injections for knee osteoarthritis: importance for individual patient outcomes. British Journal of Sports Medicine 2016;50:84-92. Available at: https://bjsm.bmj.com/content/50/2/84.full Accessed Feb 2021
- 23. American College of Rheumatology 2012 Recommendations for the Use of Nonpharmacologic and Pharmacologic Therapies in Osteoarthritis of the Hand, Hip, and Knee. Arthritis Care Res, 64: 465-474. Available at: https://doi.org/10.1002/acr.21596

- https://onlinelibrary.wiley.com/doi/full/10.1002/acr.21596 Accessed Feb 2021
- 24. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Management of Osteoarthritis of the Hand, Hip, and Knee. Arthritis Care Res, 72: 149-162. Link Accessed Feb 2021
- 25. National Institute for Health and Care Excellence (NICE). Osteoarthritis: care and management. 2014 Feb. (NICE clinical guideline; no. 177).
- 26. Osteoarthritis Research Society International (OARSI). OARSI guidelines for the non-surgical management of knee, hip, and polyarticular osteoarthritis. Osteoarthritis and Cartilage. 2019 Nov; 27(11):1578-1589. Available at: Link Accessed Feb 2021
- 27. Department of Veterans Affairs/Department of Defense (VA/DoD). VA/DoD Clinical Practice Guideline for the Non-surgical Management of Hip & Knee Osteoarthritis. Version 2.0 2020. Available at: Link Accessed Feb 2021

| SUMMARY OF REVIEW/REVISIONS          | DATE                       |
|--------------------------------------|----------------------------|
| REVISION- Notable revisions:         | Q4 2022                    |
| Required Medical Information         |                            |
| Continuation of Therapy              |                            |
| Prescriber Requirements              |                            |
| Age Restrictions                     |                            |
| Quantity                             |                            |
| Compendial Approved Off-Labeled Uses |                            |
| References                           |                            |
| Q2 2022 Established tracking in new  | Historical changes on file |
| format                               | -                          |
|                                      |                            |