

Effective Date: 07/28/2021 Last P&T Approval/Version: 07/28/2021 Next Review Due By: 07/1/2022

Policy Number: C21454-A

Topical Acne and Topical Rosacea Illinois Medicaid Only

PRODUCTS AFFECTED

Azelex CREA 20% (Azelaic Acid Cream 20%), Cleocin-T GEL 1% (Clindamycin Phosphate Gel 1%), Cleocin-T LOTN 1% (Clindamycin Phosphate Lotion 1%), Clinda in ETA RT 1% (Clindamycin Phosphate Swab 1% & Cleanser Kit), Clindacin ETZ SWAB 1% (Clindamycin Phosphate Swab 1%), Clindamycin Phosphate GEL 1%, Clindamycin Phosphate LOTN 1%, Clindamycin Phosphate SOLN 1%, Clindamycin Phosphate FOAM 1%, Clindagel GEL 1% (Clindamycin Phosphate Gel 1%), Clindamycin Phosphate SWAB 1%, Clindacin-SWAB 1% (Clindamycin Phosphate Swab 1%), Evoclin FOAM 1% (Clindamycin Phosphate Foam 1%), Dapsone GEL 5% (Dapsone Gel 5%), Aczone GEL 5% (Dapsone Gel 5%), Aczone GEL 7.5% (Dapsone Gel 7.5%), Dapsone GEL 7.5% (Dapsone Gel 7.5%), Erythromycin SOLN 2% (Erythromycin Soln 2%), Erygel GEL 2% (Erythromycin Gel 2%), Erythromycin GEL 2% (Erythromycin Pads 2%), Erythromycin PADS 2% (Erythromycin Pads 2%), Klaron LOTN 10% (Sulfacetamide Sodium Lotion 10% (Acne)), Sulfacetamide Sodium (Acne) LOTN 10%, Amzeeq FOAM 4% (Minocycline HCl Micronized Foam 4%), Adapalene-Benzoyl Peroxide GEL 0.1-2.5% (Adapalene-Benzoyl Peroxide Gel 0.1-2.5%), Epiduo Forte GEL 0.3-2.5% (Adapalene-Benzoyl Peroxide Gel 0.3-2.5%), Aktipak PACK 5-3% (Benzoyl Peroxide-Erythromycin Gel Pack 5-3%), Benzamycin GEL 5-3% (Benzoyl Peroxide-Erythromycin Gel 5-3%), Benzoyl Peroxide-Erythromycin GEL 5-3% (Benzoyl Peroxide-Erythromycin Gel 5-3%), BenzaClin GEL 1-5% (Clindamycin Phosphate Benzoyl Peroxide Gel 1-5%), BenzaClin with Pump GEL 1-5% (Clindamycin Phosphate-Benzoyl Peroxide Gel 1-5%), Clindamycin Phosphate-Benzoyl Peroxide Gel 1-5%, Clindamycin Phosphate-Benzoyl Peroxide Gel 1.2-2.5%, Acanya GEL 1.2-2.5% (Clindamycin Phosphate-Benzoyl Peroxide Gel 1.2-2.5%), Onexton GEL 1.2-3.75% (Clindamycin Phosphate-Benzoyl Peroxide Gel 1.2-3.75%), Clindamycin Phos-Benzoyl Perox GE 1.2-5%, Neuac GEL 1.2-5% (Clindamycin Phosph-Benzoyl Peroxide (Refrig) Gel 1.2 (1)-5%), Clindacin Pac KIT 1% (*Clindamycin Phosphate Swab 1% & Cleanser Kit), Clindamycin-Tretinoin GEL 1.2-0.025% (Clindamycin Phosphate-Tretinoin Gel 12-0.025%), Ziana GEL 1.2-0.025% (Clindamycin Phosphate-Tretinoin Gel 1.2-0.025%), Sulfacetamide Sodium-Sulfur LIQD 9-4% (Sulfacetamide Sodium w/ Sulfur Wash 9-4%), Sumaxin Wash LIQD 9-4% (Sulfacetamide Sodium w/ Sulfur Wash 9-4%), Sulfacetamide Sodium-Sulfur LIQD 9-4.5% (Sulfacetamide Sodium w/ Sulfur Wash 9-4.5%), Sumadan Wash LIQD 9-4.5% (Sulfacetamide Sodium w/ Sulfur Wash 9-4.5%), Sulfacetamide Sodium-Sulfur LIQD 9.8-4.8% (Sulfacetamide Sodium w/ Sulfur Cleanser 9.8-4.8%), Sulfacetamide Sodium-Sulfur LIQD 10-2% (Sulfacetamide Sodium w/ Sulfur Cleanser 10-2%), BP 10-1 EMUL 10-1% (Sulfacetamide Sodium w/ Sulfur Emulsion 10-1%), Sulfacetamide Sodium w/ Sulfur Emulsion 10-5%, Sulfacetamide Sodium w/ Sulfur Susp 8-4%, Sumaxin TS SUSP 8-4% (Sulfacetamide Sodium w/ Sulfur Susp 8-4%), Sulfacetamide Sodium w/ Sulfur Cream 10-2%, SSS 10-5 CREA 10-5% (Sulfacetamide Sodium w/ Sulfur Cream 10-5%), Sulfacetamide Sodium w/ Sulfur Cream 10-5%, SSS 10-5 FOAM 10-5% (Sulfacetamide Sodium w/ Sulfur Foam 10-5%), Sulfacetamide Sodium w/

Sulfur Cleansing Pad 10-4%, Sumaxin PADS 10-4% (Sulfacetamide Sodium w/ Sulfur Cleansing Pad 10-4%), BP Cleansing Wash EMUL 10-4% (Sulfacetamide Sodium-Sulfur in Urea Emulsion 10-4%). Sulfacetamide Sodium-Sulfur in Urea Emulsion 10-5%. Sumadan KIT 9-4.5% (Sulfacetamide Sod-Sulfur Wash 9-4.5% & Skin Cleanser Kit), Sumaxin CP KIT 10-4% (Sulfacetamide Sod-Sulfur Pad 10-4% & Skin Cleanser Kit), Sumadan XLT KIT 9-4.5% (Sulfacetamide Sod-Sulfur Wash 9-4.5% & Sunscreen Kit), Neuac KIT 1.2-5% (Clindamycin-Benzoyl Perox Gel 1.2-5% & Moisturizer Cr Kit); Finacea FOAM 15% (Azelaic Acid Foam 15%), Azelaic Acid GEL 15%, Finacea GEL 15% (Azelaic Acid Gel 15%), GEL 0.33% (Brimonidine Tartrate Gel 0.33% (Base Equivalent)), Soolanta CREAM 1% (Ivermectin Cream 1%, Ivermectin CREAM 1%, metroNIDAZOLE CREAM 0.75% (Metronidazole Cream 0.75%), MetroCream CREAM 0.75%, Rosadan CREAM 0.75% (Metronidazole Cream 0.75%), Noritate CREA 1% (Metronidazole Cream 1%) metroNIDAZOLE GEL 0.75%, Rosadan GEL 0.75% (Metronidazole Gel 0.75%), Metrogel GEL 1% (Metronidazole Gel 1%), metroNIDAZOLE GEL 1%, metroNIDAZOLE 0.75%, Rosadan KIT 0.75% (Gel) (Metronidazole Gel 0.75% w/ Cleanser Kit), Rosadan KIT 0.75% (Cream) (Metronidazole Cream 0.75% w/ Cleanser Kit), Zilxi FOAM 1.5% (Minocycline HCI Micronized Foam 1.5%), Rhofade CREA 1% (Oxymetazoline HCI Cream 1%)

COVERAGE POLICY

Coverage for services, procedures, medical devices, and drugs are dependent upon benefit eligibility as outlined in the member's specific benefit plan. This Coverage Guideline must be read in its entirety to determine coverage eligibility, if any.

This Coverage Guideline provides information related to coverage determinations only and does not imply that a service or treatment is clinically appropriate or inappropriate. The provider and the member are responsible for all decisions regarding the appropriateness of care. Providers should provide Molina Healthcare complete medical rationale when requesting any exceptions to these guidelines

Documentation Requirements:

Molina Healthcare reserves the right to require that additional documentation be made available as part of its coverage determination; quality improvement; and fraud; waste and abuse prevention processes. Documentation required may include, but is not limited to, patient records, test results and credentials of the provider ordering or performing a drug or service. Molina Healthcare may deny reimbursement or take additional appropriate action if the documentation provided does not support the initial determination that the drugs or services were medically necessary, not investigational or experimental, and otherwise within the scope of benefits afforded to the member, and/or the documentation demonstrates a pattern of billing or other practice that is inappropriate or excessive

DIAGNOSIS:

Açne Vulgaris, Rosacea

REQUIRED MEDICAL INFORMATION:

- A. ACNE VULGARIS
 - Documentation of a diagnosis of acne vulgaris AND
 - For Non-preferred PDL products (per Illinois Medicaid Preferred Drug List):
 Documentation of an adequate trial (of at least 4 weeks) of a topical antibiotic
 AND one of the following, used in combination: topical retinoid or topical benzoyl peroxide.

Note: Combination therapy may be achieved using single agent products or combination products.

Molina Reviewer Note: Preferred topical acne products require prior authorization for members under the age of 10 or greater than 21 per Illinois HFS. Trial and failure of other agents not required if the product is preferred. AND

- 3. Prescriber attests to (or the clinical reviewer has found that) the member not having any FDA labeled contraindications that haven't been addressed by the prescriber within the documentation submitted for review [Topical clindamycin containing products are contraindicated in individuals with a history of hypersensitivity to preparation containing clindamycin or lincomycin, a history of regional enteritis or ulcerative colitis, or a history of antibiotic-associated colitis Topical erythromycin is contraindicated in those individuals who have shown hypersensitivity to any of its components; topical benzoyl peroxide containing products are contraindicated in those individuals who have shown hypersensitivity to any of its components; Topical Sodium sulfacetamide/sulfur products are contraindicated in patients having known hypersensitivity to sulfonamides, sulfur or any other component of this preparation and those patients with kidney disease; Topical sulfacetamide products are contraindicated for use by patients having known hypersensitivity to sulfonamides or any other component of this preparation; Topical minocycline products are contraindicated in persons who have shown hypersensitivity to any of the tetracyclines or any of the ingredients of the product.]
- 4. For members under the age of 10, documentation that the requested product is FDA labeled or compendia supported for the member's current age AND
- 5. IF THIS IS A NON-FORMULARY PRODUCT(NOT LISTED ON THE ILLINOIS MEDICAID PREFERRED DRUG LIST): Documentation of trial/failure of or intolerance to a majority (not more than 3) of the formulary (PDL) alternatives for the given diagnosis. If yes, please submit documentation including medication(s) tried, dates of trial(s) and reason for treatment failure(s).

B. ROSACEA

- Documentation of a diagnosis of rosacea AND
- 2. For non-preferred PDL products (per Illinois Medicaid Preferred Drug List):

 Documentation of a trial and failure of a formulary preferred topical metronidazole product

Molina Reviewer Note: Preferred topical acne products, which may also be FDA labeled for rosacea, require prior authorization for members under the age of 10 or greater than 21 per Illinois HFS. Trial and failure of other agents not required if the product is preferred.

AND

3. Prescriber attests to (or the clinical reviewer has found that) the member not having any FDA labeled contraindications that haven't been addressed by the prescriber within the documentation submitted for review [Topical Sodium sulfacetamide/sulfur products are contraindicated in patients having known hypersensitivity to sulfonamides, sulfur or any other component of this preparation and those patients with kidney disease; Topical sulfacetamide products are contraindicated for use by patients having known hypersensitivity to sulfonamides or any other component of this preparation; Topical minocycline

products are contraindicated in persons who have shown hypersensitivity to any of the tetracyclines or any of the ingredients of the product; Finacea Gel is contraindicated in individuals with a history of hypersensitivity to propylene glycol or any other component of the formulation; Topical metronidazole products are contraindicated in individuals with a history of hypersensitivity to metronidazole, or other ingredients of the formulations.]

4. IF THIS IS A NON-FORMULARY PRODUCT (NOT LISTED ON THE ILLINOIS MEDICAID PREFERRED DRUG LIST): Documentation of trial/failure of or intolerance to a majority (not more than 3) of the formulary (PDL) alternatives for the given diagnosis. If yes, please submit documentation including medication(s) tried, dates of trial(s) and reason for treatment failure(s).

CONTINUATION OF THERAPY:

A. ALL INDICATIONS:

- Adherence to therapy at least 85% of the time as verified by the prescriber or member medication fill history
- Prescriber attestation that member has had no intolerable adverse effects or drug toxicity AND
- 3. Documentation of positive clinical response as demonstrated by improvements in the condition's signs and symptoms

DURATION OF APPROVAL:

Initial authorization: 12 months, Continuation of Therapy: 12 months

PRESCRIBER REQUIREMENTS

None

AGE RESTRICTIONS (for Non-preferred products only):

ACNE:

Note: Preferred topical acre products require prior authorization for members under the age of 10 or

greater than 21 per Illinois HFS Aczone Gel 7.5%: Age 9 and older

Amzeeq: Age 9 and older

Epiduo 0.1%;2.5%: Age 9 and older

Other Topical Acne Agents: Age 10 or older

ROSACEA:

Note: Preferred topical acne products, which may also be FDA labeled for rosacea, require prior authorization for members under the age of 10 or greater than 21 per Illinois HFS

Mirvaso, Finacea, Soolantra, Metrogel Brand, MetroCream Brand, Noritate Brand, Zilxi: Age 18 or older Non-preferred sulfacetamide sodium containing products, Azelex cream 20%: Age 10 and older

QUANTITY:

See Illinois PDL for quantity limitations

Page 4 of 10

PLACE OF ADMINISTRATION:

The recommendation is that topical medications in this policy will be for pharmacy benefit coverage and patient self-administered.

DRUG INFORMATION

ROUTE OF ADMINISTRATION:

Topical

DRUG CLASS:

Acne Combinations

FDA-APPROVED USES:

ACANYA Gel: indicated for the topical treatment of acne vulgaris in patients 12 years or older

Aczone Gel 7.5%: indicated for the topical treatment of acne vulgaris in patients 9 years of age and older

Amzeeq foam: indicated to treat inflammatory lesions of non-nodular moderate to severe acne vulgaris in patients 9 years of age and older.

Azelex: indicated for the topical treatment of mild-to-moderate inflammatory acne vulgaris

BP 10-1(sodium sulfacetamide 10% and sulfur 1%): indicated in the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis.

Clindacin P: indicated in the treatment of acne vulgaris

Cleocin T topical solution, Cleocin T Topical Gel, Cleocin T Topical Lotion: indication in the treatment of acne vulgaris

EPIDUO FORTE: indicated for the topical treatment of acne vulgaris.

ERYGEL Topical Gel: indicated for the topical treatment of acne vulgaris.

EVOCLIN Foam; indicated for acne vulgaris in patients 12 years and older.

FINACEA (azelaic acid) foam, gel: indicated for the topical treatment of inflammatory papules and pustules of mild to moderate rosacea

Klaron Lotion: indicated in the topical treatment of acne vulgaris.

Metrogel 1%, MetroCream 0.75%, Noritate 1% cream: indicated for the treatment of inflammatory papules, pustules, and erythema of rosacea

MIRVASO (brimonidine) topical gel: indicated for the topical treatment of persistent (non-transient) facial erythema of rosacea in adults 18 years of age or older

Neuac Gel, 1.2%/5%:indicated for the topical treatment of inflammatory acne

ONEXTON Gel: indicated for the topical treatment of acne vulgaris in patients

Drug and Biologic Coverage Criteria 12 years of age and older.

RHOFADE (oxymetazoline hydrochloride) cream, 1%: indicated for the topical treatment of persistent facial erythema associated with rosacea in adults.

Rosadan 0.75% Cream and Gel: indicated for topical application in the treatment of inflammatory papules and pustules of rosacea.

SOOLANTRA (ivermectin) cream, 1%: s indicated for the treatment of inflammatory lesions of rosacea

Sumaxin (sodium sulfacetamide 10% & sulfur 4%) Cleansing Pads: indicated in the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis

Sumadan (sodium sulfacetamide 9% & sulfur 4.5%) Wash: indicated for the topical control of acne vulgaris, acne rosacea and seborrheic dermatitis

ZIANA (clindamycin phosphate 1.2% and tretinoin 0.025%) Gel: indicated for the topical treatment of acne vulgaris in patients 12 years or older

ZILXI (minocycline) topical foam: indicated for the treatment of inflammatory lesions of rosacea in adults

COMPENDIAL APPROVED OFF-LABELED USES

UpToDate, Micromedex, AHFS or Clinical Pharmacology

APPENDIX

APPENDIX:

Treatment algorithm for the management of acne vulgaris in adolescents and young adults. Adopted from the 'Guidelines of care for the management of acne vulgaris' J Am Acad Dermatol 2016

•			
	Mild	Moderate	Severe
1st Line Treatment	Benzoyl Peroxide (BP) or Topical Retinoid -or- Topical Combination Therapy** BP + Antibiotic or Retinoid + BP or Retinoid + BP + Antibiotic	Topical Combination Therapy** BP + Antibiotic or Retinoid + BP or Retinoid + BP + Antibiotic -or- Oral Antibiotic + Topical Retinoid + BP -or- Oral Antibiotic + Topical Retinoid + BP + Topical Antibiotic	Oral Antibiotic + Topical Combination Therapy** BP + Antibiotic or Retinoid + BP or Retinoid + BP + Antibiotic -or- Oral Isotretinoin
Alternative Treatment	Add Topical Retinoid or BP (if not on already) -or- Consider Alternate Retinoid -or- Consider Topical Dapsone	Consider Alternate Combination Therapy -or- or- Consider Change in Oral Antibiotic -or- Add Combined Oral Contraceptive or Oral Spironolactone (Females) -or- Consider Oral Isotretinoin	Consider Change in Oral Antibiotic -or- Add Combined Oral Contraceptive or Oral Spironolactone (Females) -or- Consider Oral Isotretinoin

Fig. 1. Treatment algorithm for the management of acne vulgaris in adolescents and young adults. The double asterists (**) indicate that the drug may be prescribed as a fixed combination product or as separate component. BP. Benzoyl peroxide.

BACKGROUND AND OTHER CONSIDERATIONS

BACKGROUND:

None

CONTRAINDICATIONS/EXCLUSIONS/DISCONTINUATION:

All other uses of topical acne products are considered experimental/investigational and therefore, will follow Molina's Off-Label policy.

OTHER SPECIAL CONSIDERATIONS:

Isotretinoin have a separate policy for alternative acne treatments. Single agent Topical Retinoids to be reviewed with Global Formulary Exception Criteria (NF, QL, ST, etc.) Policy Number: C10169-A.

CODING/BILLING INFORMATION

Note: 1) This list of codes may not be all-inclusive. 2) Deleted codes and codes which are not effective at the time the service is rendered may not be eligible for reimbursement

HCPCS CODE	DESCRIPTION
NA	

AVAILABLE DOSAGE FORMS:

Cleocin-T GEL 1% (Clindamycin Phosphate Gel 1%)

Cleocin-T LOTN 1% (Clindamycin Phosphate Lotion 1%)

Clindacin ETZ KIT 1% (*Clindamycin Phosphate Swab 1% & Cleanser Kit***)

Clindacin ETZ SWAB 1% (Clindamycin Phosphate Swab 1%)

Clindamycin Phosphate GEL 1%

Clindamycin Phosphate LOTN 1%

Clindamycin Phosphate SOLN 1%

Clindamycin Phosphate FOAM 1%

Clindagel GEL 1% (Clindamycin Phosphate Gel 1%)

Clindamycin Phosphate SWAB 1%

Clindacin-P SWAB 1% (Clindamycin Phosphate Swab 1%)

Evoclin FOAM 1% (Clindamycin Phosphate Foam 1%)

Dapsone GEL 5% (Dapsone Gel 5%)

Aczone GEL 5% (Dapsone Gel 5%)

Aczone GEL 7.5% (Dapsone Gel 7.5%)

Dapsone GEL 7.5% (Dapsone Gel 7.5%)

Erythromycin SOLN 2% (Erythromycin Soln 2%)

Erygel GEL 2% (Erythromycin Gel 2%)

Erythromycin GEL 2% (Erythromycin Gel 2%)

Ery PADS 2% (Erythromycin Pads 2%)

Erythromycin PADS 2% (Erythromycin Pads 2%)

Finacea (azelaic acid) foam, gel

Klaron LOTN 10% (Sulfacetamide Sodium Lotion 10% (Acne))

Sulfacetamide Sodium (Acne) LOTN 10%

Amzeeq FOAM 4% (Minocycline HCl Micronized Foam 4%)

Adapalene-Benzoyl Peroxide GEL 0.1-2.5% (Adapalene-Benzoyl Peroxide Gel 0.1-2.5%)

Epiduo Forte GEL 0.3-2.5% (Adapalene-Benzoyl Peroxide Gel 0.3-2.5%)

Aktipak PACK 5-3% (Benzoyl Peroxide-Erythromycin Gel Pack 5-3%)

Benzamycin GEL 5-3% (Benzoyl Peroxide-Erythromycin Gel 5-3%)

Benzoyl Peroxide-Erythromycin GEL 5-3% (Benzoyl Peroxide-Erythromycin Gel 5-3%)

BenzaClin GEL 1-5% (Clindamycin Phosphate-Benzoyl Peroxide Gel 1-5%)

BenzaClin with Pump GEL 1-5% (Clindamycin Phosphate-Benzoyl Peroxide Gel 1-5%)

Clindamycin Phosphate-Benzoyl Peroxide Gel 1-5%

Clindamycin Phosphate-Benzoyl Peroxide Gel 1.2-2.5%

Acanya GEL 1.2-2.5% (Clindamycin Phosphate-Benzoyl Peroxide Gel 1.2-2.5%)

Onexton GEL 1.2-3.75% (Clindamycin Phosphate-Benzoyl Peroxide Gel 12-3.75%)

Clindamycin Phos-Benzoyl Perox GEL 1.2-5%

Neuac GEL 1.2-5% (Clindamycin Phosph-Benzoyl Peroxide (Refrig) Gel 1.2 (1)-5%)

Clindacin Pac KIT 1% (*Clindamycin Phosphate Swab 1% & Cleanser Kit)

Clindamycin-Tretinoin GEL 1.2-0.025% (Clindamycin Phosphate-Tretinoin Gel 1.2-0.025%)

Ziana GEL 1.2-0.025% (Clindamycin Phosphate-Tretinoin Gel 1.2-0.025%)

Sulfacetamide Sodium-Sulfur LIQD 9-4% (Sulfacetamide Sodium W/Sulfur Wash 9-4%)

Sumaxin Wash LIQD 9-4% (Sulfacetamide Sodium w/ Sulfur Wash 9-4%)

Sulfacetamide Sodium-Sulfur LIQD 9-4.5% (Sulfacetamide Sodium w/ Sulfur Wash 9-4.5%)

Sumadan Wash LIQD 9-4.5% (Sulfacetamide Sodium w/ Sulfur Wash 9-4.5%)

Sulfacetamide Sodium-Sulfur LIQD 9.8-4.8% (Sulfacetamide Sodium w/ Sulfur Cleanser 9.8-4.8%)

Sulfacetamide Sodium-Sulfur LIQD 10-2% (Sulfacetamide Sodium w/ Sulfur Cleanser 10-2%)

BP 10-1 EMUL 10-1% (Sulfacetamide Sodium w/ Sulfur Emulsion 10-1%)

Sulfacetamide Sodium w/ Sulfur Emulsion 10-5%

Sulfacetamide Sodium w/ Sulfur Susp 8-4%

Sumaxin TS SUSP 8-4% (Sulfacetamide Sodium w/ Sulfur Susp 8-4%)

Sulfacetamide Sodium w/ Sulfur Cream 10-2%

SSS 10-5 CREA 10-5% (Sulfacetamide Sodium w/ Sulfur Cream 10-5%)

Sulfacetamide Sodium w/ Sulfur Cream 10-5%

SSS 10-5 FOAM 10-5% (Sulfacetamide Sodium w/ Sulfur Foam 10-5%)

Sulfacetamide Sodium w/ Sulfur Cleansing Pad 10-4%

Sumaxin PADS 10-4% (Sulfacetamide Sodium w/ Sulfur Cleansing Pad 10-4%)

BP Cleansing Wash EMUL 10-4% (Sulfacetamide Sodium-Sulfur in Urea Emulsion 10-4%)

Sulfacetamide Sodium-Sulfur in Urea Emulsion 10-5%

Sumadan KIT 9-4.5% (Sulfacetamide Sod-Sulfur Wash 9-4.5% & Skin Cleanser Kit)

Sumaxin CP KIT 10-4% (Sulfacetamide Sod-Sulfur Pad 10-4% & Skin Cleanser Kit)

Sumadan XLT KIT 9-4.5% (Sulfacetamide Sod-Sulfur Wash 9-4.5% & Sunscreen Kit)

Neuac KIT 1.2-5% (Clindamycin-Benzoyl Perox Gel 1.2-5% & Moisturizer Cr Kit)

MIRVASO (brimonidine) topical gel 0.33%

SOOLANTRA (ivermectin) cream, 1%

Rosadan KIT 0.75% (Gel)

Rosadan KIT 0.75% (Cream)

MetroCream CREA 0.75%

Noritate CREA 1%

Metrogel GEL 1% Zilxi FOAM 1.5% Rhofade CREA 1%

REFERENCES

- Cleocin T (clindamycin phosphate topical solution, gel, lotion) [prescribing information]. New York, NY: Pharmacia & Upjohn Co. Division of Pfizer; December 2019
- Zaenglein, Andrea L. et al. Guidelines of care for the management of acne vulgaris. J
 Am AcadDermatol. 2016 May;74(5):945-73.e33. doi: 10.1016/j.jaad.2015.12.037. Epub 2016
 Feb 17.Available at: https://www.jaad.org/article/S0190-9622(15)02614-6/fulli.xii...45-essed Feb 2020
- 3. Clindagel (clindamycin phosphate) topical gel [prescribing information]. San Antonio, Texas: Valeant Pharmaceuticals North America LLC: November 2017
- 4. Evoclin (clindamycin phosphate) foam 1% [prescribing information]. Research Triangle Park, NC: Manufactured for: Stiefel Laboratories, Inc.; January 2012
- Cleocin P (clindamycin phosphate topical solution) [prescribing information]. Fairfield, NJ: Medimetriks Pharmaceuticals, Inc.; November 2016
- 6. ACZONE (dapsone) Gel [prescribing information]. Irvine, Ca. Allergan; February 2016
- 7. Erygel Topical Gel [prescribing information]. Morgantown, WV: Mylan Pharmaceuticals Inc.; June 2018
- KLARON (sodium sulfacetamide lotion) Lotion, 10% [prescribing information]; Bridgewater, NJ;
 Dermik Laboratories, a business of sanofi-aventis U.S. LLC
- 9. Epiduo Forte [prescribing information]; For Worth TX: Galderma Laboratories, L.P.: July 2015
- 10. BenzaClin Topical Gel (clindamycin benzoyl peroxide gel) [prescribing information]; Bridgewater, NJ; Dermik Laboratories, a business of sanofi-aventis U.S. LLC: June 2010
- 11. ONEXTON (clindamycin phosphate and benzoyl peroxide) gel [prescribing information]; Bridgewater, NJ: Bausch Health US, LLC. April 2020
- 12. Acanya (clindamycin phosphate and benzoyl peroxide) gel [prescribing information]; Bridgewater, NJ: Bausch Health US, LLC. September 2020
- 13. Sumaxin (sodium sulfacetamide 10% and sulfur 4%) [prescribing information]; Fairfield, NJ, Medimetriks Pharmaceuticals, Inc; October 2017
- 14. Sumadan(sodium sulfacetamide 10% and sulfur 4%) Wash [prescribing information]; Fairfield, NJ, Medimetriks Pharmaceuticals, Inc; March 2013
- 15. BP 10-1 (sodum sulfacetamide 10% and sulfur 15) [prescribing information]; Alpharetta, GA: Acella Pharmaceuticals, LLC: March 2019
- 16. Neuac (clindamycin phosphate and benzoyl peroxide gel [prescribing information]; Fairfield, NJ: Medimetriks Pharmaceuticals, Inc.; November 2015
- 17. ZIANA (clindamycin phosphate 1.2% and tretinoin 0.025%) Gel [prescribing information]; Buffalo, NY: Medicis, The Dermatology Company; November 2006
- 18 Amzeeg (minocycline) topical foam [prescribing information]; Bridgewater, NJ: VYNE Pharmaceuticals Inc.; January 2021
- 19. FINACEA (azelaic acid) foam [prescribing information]; Madison, NJ: LEO Pharma Inc.; December 2020
- 20. FINACEA (azelaic acid) gel[prescribing information]; Pine Brook, NJ: Intendis Inc.; July2020
- 21. MIRVASO (brimonidine) topical gel [prescribing information];Fort Worth, TX, Galderma Laboratories; November 2011
- 22. SOOLANTRA (ivermectin) cream, 1% [prescribing information]; Fort Worth, TX, Galderma Laboratories; July 2018
- 23. Rosadan (Metronidazole Topical Cream, 0.75%) Cream [prescribing information]; Fairfield, NJ:

Medimetriks Pharmaceuticals, Inc.; July 2017

- 24. ZILXI (minocycline) topical foam [prescribing information]; Bridgewater, NJ: VYNE Pharmaceuticals Inc.; January 2021
- 25. RHOFADE (oxymetazoline hydrochloride) cream, 1% [prescribing information]; Charleston, SC: EPI Health; November 2019

